arXiv:cond-mat/9802284v1 [cond-mat.stat-mech] 26 Feb 1998

Non-perturbative renormalization of the KPZ growth dynamics

C. Castellano[(1)], M. Marsili[(2)] and L. Pietronero[(1,3)]
[(1)] *The "Abdus Salam" I.C.T.P. Strada Costiera 11, I-34100 Trieste*
[(2)] *International School for Advanced Studies (SISSA), via Beirut 2-4, Trieste I-34014*
[(3)] *Dipartimento di Fisica and INFM Unit, University of Rome "La Sapienza", I-00185 Roma*

We introduce a nonperturbative renormalization approach which identifies stable fixed points in any dimension for the Kardar-Parisi-Zhang dynamics of rough surfaces. The usual limitations of real space methods to deal with anisotropic (self-affine) scaling are overcome with an indirect functional renormalization. The roughness exponent α is computed for dimensions $d = 1$ to 8 and it results to be in very good agreement with the available simulations. No evidence is found for an upper critical dimension. We discuss how the present approach can be extended to other self-affine problems.

PACS numbers: 05.40.+j,64.60.Ak,05.70.Ln,68.35.Fx

Several models for the growth of rough surfaces appear to belong to the universality class of the Kardar-Parisi-Zhang (KPZ) equation [1,2]:

$$\frac{\partial h(x,t)}{\partial t} = \nu \nabla^2 h + \frac{\lambda}{2}(\nabla h)^2 + \eta(x,t). \qquad (1)$$

Here the first term on the r.h.s. corresponds to surface tension, the second describes lateral growth [1] and η is white noise. The large scale properties of growing surfaces are described by two exponents: the roughness exponent α, defined by the typical height fluctuation $\delta h \sim L^\alpha$ over a linear distance L, and the dynamic exponent z [2]. For the KPZ equation, the exponents satisfy the scaling relation $z + \alpha = 2$ as a consequence of Galileian invariance in the related Burgers equation [3]. It is then sufficient to concentrate attention on α. In the usual field theory approach a perturbation theory is defined with respect to the non-linear term (λ). The corresponding renormalization group (RG) reveals [2] that the physics of the KPZ is related to a strong coupling fixed point, which is unaccessible by perturbative methods. Except for $d = 1$, where, thanks to a fluctuation-dissipation theorem, an analytic solution is possible, the problem appears of strong coupling nature and not much can be done from this theoretical perspective.

Extensive numerical simulations have been carried out for restricted solid on solid (RSOS) models [4], for $d = 1$ to 7. These are discrete models which belong to the KPZ universality class [5] and are optimal for numerical simulations [2,4]. These studies show a gradual decrease of the value of α, without evidence for an upper critical dimension. On the other hand, several analytical approaches [6] suggest an upper critical dimension $d_c = 4$ above which one should have $\alpha = 0$.

In strong coupling situations, real space methods perform sometimes much better than continuum field theory. This is particularly true for self-organized non-equilibrium systems such as fractal growth [7], sandpile models [8] and models with quenched disorder [9] where real space methods give quite accurate results. However, rough surfaces show anisotropic scaling properties (self-affinity) and, in such a case, it is difficult to define the "internal degrees of freedom" for the block transformation in real space. Here we overcome this basic problem by defining the renormalization transformation in an indirect way that permits to integrate over the internal degrees of freedom without the necessity of specifying them explicitly. This approach is rather general and it should be useful also for other self-affine structures.

Our starting point is to introduce a parametrization of the dynamics at a generic scale. Then we devise a RG where a scale transformation defines a flux in the parameter space of possible dynamics. The fixed point of the renormalization transformation defines the *scale invariant dynamics* and it describes the asymptotic scaling properties of the system. The key point lies in the choice of a correct parametrization of the dynamics which captures the physics of the problem but is also simple enough to allow for explicit calculations. The main feature of the KPZ dynamics is lateral growth [1,5]. This allows us to concentrate on just one renormalization parameter which is the ratio between lateral and vertical growth rates. At the simplest level the method is fully analytical while for accurate results or $d > 1$ we resort to the Monte Carlo method to compute the renormalization functions. We obtain stable and attractive fixed points for any dimension from $d = 1$ to 8. From them we compute the roughness exponent α which results to be in excellent agreement with the available simulations [4] up to $d = 7$. From the values of α and the stability of the fixed points we conclude that there is no upper critical dimension, at least up to $d = 8$.

Let us consider a rough surface and define a block transformation for its profile as indicated in Fig. 1. The coarse grained description is obtained by a partition of the $d + 1$ dimensional space in cells of lateral size $L_k = L_0 \cdot 2^k$ and thickness $h_k \simeq L_k^\alpha$. Each cell of the partition is then declared to be full or empty according to some majority rule. With respect to the usual

RG approaches one should note that the definition of the coarse-graining transformation depends explicitly on the exponent α which, on its turn, is just what we should compute in the end. In addition the shape of the block variables changes with the scale because, since in general $\alpha \neq 1$, the ratio h_k/L_k will depend on the scale k. For these reasons the development of real space methods for self-affine problems has been extremely limited.

With this coarse grained description of the interface in mind, let us now introduce the *effective* dynamics at the same level of coarse graining. The dynamics at scale k is defined in terms of transitions rates. At the microscopic scale these correspond to the possible growth processes occurring along the surface, as specified by the model definition (for example in Eden growth vertical and lateral growth have the same rates [2]). At a generic scale L_k we generalize this situation in order to allow for proliferation in the RG. We set the rate of the vertical growth nominally equal to L_k and define a new parameter, λ_k, for the lateral growth. The growth rate for a generic scale k can then be expressed as:

$$P[h(i) \to h(i)+1] \equiv L_k + \lambda_k \sum_{jnni} \max[0, h(j) - h(i)] \quad (2)$$

The term L_k is the contribution of the vertical growth and the sum over neighbor block sites j counts the contribution of lateral growth. We assume that $h(i)$ is expressed in units of h_k and we neglect overhangs [10].

We can now address the following question: Given a description of the system at scale k, defined by the parameters (L_k, h_k, λ_k) what will be corresponding properties (L_{k+1}, h_{k+1}, λ_{k+1}) at the scale $L_{k+1} = 2L_k$?

A standard answer to this question, in the real space RG framework, is to partition a configuration at a given scale into sub-configurations at a finer scale of coarse graining (thereby integrating over degrees of freedom). This method is problematic because of the self-affine nature of the problem. A different method, which we introduce here, is to consider a system of size $L = L_{k+2}$ and its descriptions at two different scales, say $L/2 = L_{k+1}$ and $L/4 = L_k$. Imposing that the computation of the same quantity in the two different descriptions gives the same result, leads to a RG equation relating the dynamics at scale $k+1$ to that at scale k. The natural quantity to consider in the case of surfaces is the roughness which, for a system of size L, is given by

$$W^2(L) = \frac{1}{L^d} \sum_i [h(i) - \bar{h}]^2 \quad (3)$$

where $\bar{h}$ is the average height. This has the nice property, in the perspective of the RG calculation, that if one partitions the system in cells of size b

$$W^2(L) = W^2(L/b) + W^2(b) \quad (4)$$

i.e. the roughness of the system is equal to the average of the roughness in the individual cells, plus the roughness of the average height of the various cells. Our strategy will then be to compute the roughness $W^2_{k+2} = W^2(L_{k+2})$ over a linear size L_{k+2} first using the description at scale $k+1$ and then at scale k.

In order to do this we can first consider a system of two cells of size L_{k+1} and then a system of four cells at size k. For simplicity we make the assumption of RSOS also for the coarse grained cells. This is actually an approximation because, upon renormalization, a dynamics which is of the restricted type ($|\Delta h| \leq 1$) at the smallest scale can easily proliferate into an unrestricted one. However with this restriction, the number of possible surface configurations is very small and an analytic treatment is possible. For the system of two sites we have only two configurations, that we classify by their height (h_1,h_2). The first one is the flat one (1,1), while the second has one step (1,2). With periodic boundary conditions and the definition (Eq. (2)) of the dynamics at scale k, the transition rates for the system of two configurations are $P_{1\to 2} = 2L_k$ and $P_{2\to 1} = L_k + 2\lambda_k$. Indeed in $P_{1\to 2}$ only vertical growth is possible (in both cells) whereas in configuration 2 only one site can grow and it has two lateral contributions. The master equation for the probability ϱ_i of configuration i is

$$\frac{\partial \varrho_i}{\partial t} = \sum_j \varrho_j P_{j\to i} - \varrho_i \sum_j P_{i\to j} \quad (5)$$

and its stationary solution is $\varrho_2 = 1 - \varrho_1 = 2/(3 + 2x_k)$, where $x_k = \lambda_k/L_k$. Using eq. (4), we can now compute the fluctuations at scale $k+1$

$$W^2_{k+1} = W^2_k + W^2(2, x_k) = W^2_k + \varrho_2(x_k) h^2_k/4. \quad (6)$$

Here W^2_k is the fluctuation at scale L_k which is present in both configurations and $W^2(2, x_k)$ is the roughness corresponding to the fluctuation of the average height at scale L_k. Clearly the flat configuration does not contribute and the roughness of the second configuration is $h^2_k/4$. We can relate W^2_k to h^2_k by observing that at scale k the essential information we retain on the height distribution is only the discretization scale h_k. This is equivalent to assuming that the distribution of the height fluctuations at this scale is a sum of two delta peaks

$$P(\Delta h) = \frac{1}{2}\delta\left(\Delta h - \frac{h_k}{2}\right) + \frac{1}{2}\delta\left(\Delta h + \frac{h_k}{2}\right) \quad (7)$$

It follows that $W^2_k = h^2_k/4$ and

$$\frac{W^2_{k+1}}{W^2_k} \equiv F_1(x_k) = 1 + \frac{2}{3 + 2x_k}. \quad (8)$$

This, at the moment, is a purely formal relation since x_k is unknown. Clearly eq. (8) also relates W^2_{k+2}/W^2_{k+1} to x_{k+1}, and combining these two relations we get:

$$\frac{W^2_{k+2}}{W^2_k} = \frac{W^2_{k+2}}{W^2_{k+1}} \cdot \frac{W^2_{k+1}}{W^2_k} = F_1(x_k)F_1(x_{k+1}). \qquad (9)$$

The roughness W^2_{k+2} can also be computed directly using the dynamics at scale k. For this we need to consider a system of four sites, at scale k. There are, always within the RSOS approximation and with periodic boundary conditions, six possible configurations of (h_1, h_2, h_3, h_4). The transition rates $P_{i\to j}$ are computed using eq. (2) and the stationary solution ϱ_i of the master equation (5) is easily found. This allows one to compute the roughness W^2_{k+2} for the 4 cells system following exactly the same procedure used for the 2 cells system, obtaining

$$\frac{W^2_{k+2}}{W^2_k} = F_2(x_k) = 1 + \frac{51 + 86x_k + 40x_k^2}{47 + 106x_k + 68x_k^2 + 8x_k^3} \qquad (10)$$

Consistency of the two descriptions requires that Eqs. (9) and (10) must yield the same quantity. This leads to our basic RG transformation for x_{k+1} as a function of x_k

$$F_1(x_{k+1}) = F_2(x_k)/F_1(x_k). \qquad (11)$$

It can also be cast into the more conventional form $x_{k+1} = R(x_k)$. A fixed point $x^* = R(x^*) \simeq 2.08779...$ exists and we can compute the roughness exponent as

$$\alpha = \frac{1}{2}\log_2\left(\frac{W^2_{k+1}}{W^2_k}\right) = \frac{1}{2}\log_2 F_1(x^*) \simeq 0.177352... \qquad (12)$$

The fixed point is also *attractive*. Close to the fixed point, we have $x_{k+1} - x^* = R(x_k) - x^* \simeq R'(x^*)(x_k - x^*)$. If $|R'(x^*)| < 1$, the distance $x_k - x^* \to 0$ as $k \to \infty$. We find $R'(x^*) = -0.03548...$ i.e. x^* is strongly attractive.

The value of the exponent is clearly too small with respect to the exact one ($\alpha = 1/2$). This is due to the RSOS approximation which, on the other hand, allows for a fully analytical treatment. The point is that even if at the microscopic scale the dynamics of the surface is of restricted type (RSOS), the effective dynamics for the block variables at a generic scale k defined by the renormalization procedure can proliferate in a non restricted dynamics. Allowing larger steps makes the problem analytically intractable. Still the calculation of the functions F_1 and F_2 can be done using a Monte Carlo method: For each value of the parameter $x_k = \lambda_k/L_k$, the growth process in cells of sizes $b = 2$ and 4 described by eq. (2), is numerically simulated. $F_1(x)$ and $F_2(x)$ are then computed as time averages in the stationary state from the roughness $W^2(b)$. In this way one sees that, allowing for larger steps, the value of α approaches the exact value $\alpha = 0.5$ with great accuracy. Convergence is reached when steps $|\Delta h| \approx 8$ are allowed. Going further, i.e. to 16, α does not change any more [11].

The Monte Carlo calculation of the functions $F_1(x)$ and $F_2(x)$ also allows to deal with higher dimensions $d > 1$. The function F_1 is then related to the roughness of a cell of size 2^d, while F_2 refers to 4^d. The method is exactly the same as for the one dimensional case. The results for $d = 1, \ldots 8$ are reported in Fig. 2. Instead of $R(x)$, Fig. 2 displays the function $\tilde{R}(\alpha)$ where $\alpha(x)$ is given in the second equality of eq. (12). This method is graphically convenient since in this way the exponents can be read from the intersection of $\tilde{R}(\alpha)$ with the line α and the stability of the fixed point can be inferred from the slope $\tilde{R}'(\alpha^*)$ of the function at the fixed point. We see that for $d = 1, \ldots, 8$ a fixed point with $\alpha > 0$ exists and it is attractive (see inset). The values of the exponent α are reported in table , together with the fixed point value $x^*(d)$ for $d = 1, \ldots, 8$. For comparison, table also reports the exponent α obtained in ref. [4] from numerical simulations up to $d = 7$. The agreement between our RG results and the simulations is rather impressive and it provides a strong support to the validity of both methods.

The extrapolation to $d \to \infty$ of $R'(x^*)$ suggests that the fixed point never becomes unstable, leading to the conclusion that no finite upper critical dimension exists for the KPZ dynamics (note that $\alpha = 0$ in our scheme would imply a stable fixed point at $x^* = \infty$). On the basis of these results, the arguments [6] leading to an upper critical dimension $d_c = 4$ should probably be reconsidered. Furthermore our results suggests a behavior of the type $\alpha(d) \sim 1/d$ and $x^*(d) \sim 2^d$ for large d. Indeed an expansion of our RG method for large dimensions, to be reported elsewhere [11], leads to such a behavior. Finally we verified that our RG scheme is stable with respect to the change of the cell sizes used. Indeed a RG function $x_{k+1} = R(x_k)$ can also be defined considering the functions $F_{\ell+1}(x) = W^2_{k+\ell+1}/W^2_k$, $F_\ell(x) = W^2_{k+\ell}/W^2_k$ and $F_1(x)$ (the case discussed above being $\ell = 1$). In $d = 1$ we found essentially the same results, whereas in larger dimensions we found the same qualitative picture with a weak dependence of the exponent on ℓ [11].

In summary we have introduced a new method to renormalize self-affine dynamics in real space. The key technical element is to effectively integrate over internal degrees of freedom in an indirect way *via* a consistency relation. This allows us to overcome the difficulty of defining a geometric fine graining procedure for self-affine structures. In some sense, the method generalizes the Fixed Scale Transformation [7] idea to self-affine structures. There are two key steps: first we introduce a description of the system and the dynamics at a generic scale L_k in terms of the parameters h_k and $\lambda_k = x_k L_k$. Secondly we derive the RG equation $x_{k+1} = R(x_k)$ with a new indirect method. The attractive fixed point x^* (self-organization) of the RG transformation identifies the *scale-invariant dynamics* and allows us to compute the exponent α.

From the point of view of the approximations involved in this method, we observe that the first step is essentially a description tool and, as such, it is in principle

exact. The approximations arise in the identification of the scale invariant dynamics, which can be made more and more accurate with the introduction of additional proliferation parameters. This explains the striking accuracy of the present method as compared to usual real space RG methods for equilibrium phenomena.

Usually real space RG methods are at their best for low dimensional systems. The topological complexity of interactions indeed increases with dimension thus requiring more and more drastic approximations for larger d. This is not the case here: Our method is exactly the same in any dimension and the computational complexity is overcome with Monte Carlo tecniques.

The nature of our approach is quite different from the usual perturbation techniques for KPZ. Indeed our parametrization of the dynamics does not include the Edwards-Wilkinson equation ($\lambda = 0$ in eq. (1)). Rather the non-interacting limit of our theory is random deposition ($\lambda = \nu = 0$).

Finally we observe that this general method can be applied to a wide variety of other dynamic critical phenomena. In this sense the present application to the KPZ problem is a very promising example.

We acknowledge interesting discussions with A. Maritan, A. Stella, C. Tebaldi and A. Vespignani.

[1] M. Kardar, G. Parisi and Y. C. Zhang, Phys. Rev. Lett. **56**, 889 (1986).
[2] T. Halpin-Healy and Y.-C. Zhang, 1995, Phys. Rep. **254**.
[3] D. Forster, D. R. Nelson and M. J. Stephen, Phys. Rev. A **16**, 732 (1977).
[4] T. Ala-Nissila, *et al*, J. Stat. Phys. **72**, 207 (1993); L.-H. Tang, *et al*, Phys. Rev. A **45**, 7162 (1992).
[5] M. Marsili, A. Maritan, F. Toigo and J. R. Banavar, Rev. Mod. Phys. **68**, 963 (1996).
[6] T. Halpin-Healy, Phys. Rev. A **42**, 711 (1990); J-P. Bouchaud and M.E. Cates, Phys. Rev. E **47**, 1455 (1993); M.A. Moore *et al*, Phys. Rev. Lett. **74**, 4257 (1995); M. Lässig and H. Kinzelbach, Phys. Rev. Lett. **78**, 906 (1997).
[7] A. Erzan, L. Pietronero and A. Vespignani, Rev. Mod. Phys. **67**, 545 (1995); R. Cafiero, A. Vespignani and L. Pietronero Phys. Rev. Lett. **70**, 3939 (1993).
[8] L. Pietronero, A. Vespignani and S. Zapperi Phys. Rev. Lett. **72**, 1690 (1994); A. Vespignani, S. Zapperi and L. Pietronero, Phys. Rev. E **51**, 1711 (1995).
[9] M. Marsili, Europhys. Lett. **28**, 385 (1994); A. Gabrielli, R. Cafiero, M. Marsili and L. Pietronero, Europhys. Lett. **38**, 491 (1997).
[10] Overhangs have been proved to be irrelevant in M. Marsili and Y.-C. Zhang, Phys. Rev. E, to appear (1998).
[11] C. Castellano, M. Marsili and L. Pietronero, preprint (1998).

FIG. 1. Coarse grained description at scale L_k of a growing interface.

FIG. 2. Renormalization group transformation $\tilde{R}(\alpha)$ for $d = 1, \ldots, 8$. The intersections with the line α yields the fixed point value $\alpha(x^*)$ of the exponent. The exponents α and the slope $\tilde{R}'(\alpha)$ at the fixed point, are plotted in the inset as a function of d. Since $|\tilde{R}'| < 1$ we conclude that the fixed points are stable.

d	1	2	3	4	5	6	7	8
x^*	0.735	2.76	7.16	15.35	31.8	62.5	125	240
$\alpha_{\rm RG}$	0.502	0.360	0.284	0.238	0.205	0.182	0.162	0.150
$\alpha_{\rm num}$	0.5	0.387	0.305	0.261	0.193	0.18	0.15	-

TABLE I. Results of the RG approach compared with numerical simulations. The first line gives the value of the fixed point critical parameter x^*. The fixed point is attractive and the corresponding exponent $\alpha_{\rm RG}$ is listed in the second line. The third line line reports the numerical results of ref. [4]. The excellent agreement agreement between our RG result and numerical simulations provides a strong support for the validity of both, and it points to the absence an upper critical dimension for this problem.


h_k
L_k


α
R'(x*)
0.5
0.0
-0.5
-1.0
0
2
4
6
8
d
0.6
0.4
0.2
0.0
α k+1
d=1
d=2
d=3
d=4
d=5
d=6
d=7
d=8
0.0
0.2
0.4
0.6
α k